

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
REQUEST FOR FILING NATIONAL PHASE OF
PCT APPLICATION UNDER 35 U.S.C. 371 AND 37 CFR 1.494 OR 1.495

09/889414

To: Hon. Commissioner of Patents
Washington, D.C. 20231



00909

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)

Atty Dkt: P 0281558 /Z 70475/UST
M# /Client Ref.

From: Pillsbury Winthrop LLP, IP Group:

Date: July 17, 2001

This is a **REQUEST** for **FILING** a PCT/USA National Phase Application based on:

International Application <u>PCT/GB00/00278</u> ↑ country code	2. International Filing Date <u>1</u> February <u>2000</u> Day MONTH Year	3. Earliest Priority Date Claimed <u>6</u> February <u>1999</u> Day MONTH Year (use item 2 if no earlier priority)
----------------------------------------------------------------------	---------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------

Measured from the earliest priority date in item 3, this PCT/USA National Phase Application Request is being filed within:

(a) ☐ 20 months from above item 3 date (b) ☒ 30 months from above item 3 date,

(c) Therefore, the due date (unextendable) is August 6, 2001

Title of Invention DRUG COMBINATIONS COMPRISING (E)-7-[4-(4-FLUOROPHENYL)-6-ISOPROPYL-2-METHYLSULFONYL] AMINO] PYRIMIDIN-5-YL] (3R,5S)-3,5-DIHYDROXYHEPT-6-ENOIC ACID AND AN INHIBITOR, INDUCER OR SUBSTRATE OF P450 ISOENZYME 3A4

Inventor(s) RAZA, Ali et al

Applicant herewith submits the following under 35 U.S.C. 371 to effect filing:

☒ Please immediately start national examination procedures (35 U.S.C. 371 (f)).

8. ☒ **A copy of the International Application** as filed (35 U.S.C. 371(c)(2)) is transmitted herewith (file if in English but, if in foreign language, file only if not transmitted to PTO by the International Bureau) including:

- a. ☒ Request;
b. ☒ Abstract;
c. 45 pgs. Spec. and Claims;
d. sheet(s) Drawing which are ☐ informal ☐ formal of size ☐ A4 ☐ 11"

9. ☒ **A copy of the International Application has been transmitted by the International Bureau.**

10. **A translation of the International Application** into English (35 U.S.C. 371(c)(2))

- a. ☐ is transmitted herewith including: (1) ☐ Request; (2) ☐ Abstract;
(3) pgs. Spec. and Claims;
(4) sheet(s) Drawing which are:
☐ informal ☐ formal of size ☐ A4 ☐ 11"
b. ☐ is not required, as the application was filed in English.
c. ☐ is not herewith, but will be filed when required by the forthcoming PTO Missing Requirements Notice per Rule 494(c) if box 4(a) is X'd or Rule 495(c) if box 4(b) is X'd.
d. ☐ Translation verification attached (not required now).

11. ☒ Please see the attached Preliminary Amendment
12. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)), i.e., **before 18th month from first priority date above in item 3, are transmitted herewith (file only if in English) including:**
13. ☒ PCT Article 19 claim amendments (if any) have been transmitted by the International Bureau
14. ☐ Translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)), i.e., of **claim amendments** made before 18th month, is attached (required by 20th month from the date in item 3 if box 4(a) above is X'd, or 30th month if box 4(b) is X'd, or else amendments will be considered canceled).
15. **A declaration of the inventor** (35 U.S.C. 371(c)(4))
a. ☐ is submitted herewith ☐ Original ☐ Facsimile/Copy
b. ☒ is not herewith, but will be filed when required by the forthcoming PTO Missing Requirements Notice per Rule 494(c) if box 4(a) is X'd or Rule 495(c) if box 4(b) is X'd.
16. **An International Search Report (ISR):**
a. Was prepared by ☒ European Patent Office ☐ Japanese Patent Office ☐ Other
b. ☒ has been transmitted by the international Bureau to PTO.
c. ☒ copy herewith (1 pg(s).) ☒ plus Annex of family members (1 pg(s).).
17. **International Preliminary Examination Report (IPER):**
a. ☒ has been transmitted (if this letter is filed after 28 months from date in item 3) in English by the International Bureau with Annexes (if any) in original language.
b. ☒ copy herewith in English.
c.1 ☐ IPER Annex(es) in original language ("Annexes" are amendments made to claims/spec/drawings during Examination) including attached amended:
c.2 ☐ Specification/claim pages # _____ claims # _____
Dwg Sheets # _____
d. ☐ Translation of Annex(es) to IPER **(required by 30th month due date, or else annexed amendments will be considered canceled).**
18. **Information Disclosure Statement** including:
a. ☒ Attached Form PTO-1449 listing documents
b. ☐ Attached copies of documents listed on Form PTO-1449
c. ☒ A concise explanation of relevance of ISR references is given in the ISR.
19. ☐ **Assignment** document and Cover Sheet for recording are attached. Please mail the recorded assignment document back to the person whose signature, name and address appear at the end of this letter.
20. ☐ Copy of Power to IA agent.
21. ☐ **Drawings** (complete only if 8d or 10a(4) not completed): _____ sheet(s) per set: ☐ 1 set informal;
☐ Formal of size ☐ A4 ☐ 11"
22. Small Entity Status ☒ is **Not** claimed ☐ is claimed (**pre-filing confirmation required**)
22(a) _____ (No.) Small Entity Statement(s) enclosed (since 9/8/00 Small Entity Statements(s) not essential to make claim)
23. **Priority** is hereby claimed under 35 U.S.C. 119/365 based on the priority claim and the certified copy, both filed in the International Application during the international stage based on the filing in (country) **GREAT BRITAIN** of:
- | | Application No. | Filing Date | | Application No. | Filing Date |
|-----|-----------------|------------------|-----|-----------------|---------------|
| (1) | 9902593.4 | February 6, 1999 | (2) | 9921063.5 | Sept. 8, 1999 |
| (3) | 9921064.3 | Sept. 8, 1999 | (4) | | |
| (5) | | | (6) | | |
- a. ☒ See Form PCT/IB/304 sent to US/DO with copy of priority documents. If copy has not been received, please proceed promptly to obtain same from the IB.
b. ☐ Copy of Form PCT/IB/304 attached.

RE: USA National Phase Filing of PCT/GB00/00278

24. Attached: 2 copies of Form PCT/IB/306

25 Per Item 17.c2, **cancel original** pages # _____, claims # _____, Drawing Sheets # _____

26. Calculation of the U.S. National Fee (35 U.S.C. 371 (c)(1)) and other fees is as follows:

Based on amended claim(s) per above item(s) ☐ 12, ☐ 14, ☐ 17, ☐ 25 (hilitte)

Total Effective Claims	minus 20 =	x \$18/\$9	= \$0	966/967
Independent Claims	minus 3 =	x \$80/\$40	= \$0	964/965
If any proper (ignore improper) Multiple Dependent claim is present,		add \$270/\$135	+0	968/969

BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(4)): →→ **BASIC FEE REQUIRED, NOW** →→→→

A. If country code letters in item 1 are **not** "US", "BR", "BB", "TT", "MX", "IL", "NZ", "IN" or "ZA"

See item 16 re:

1. Search Report was <u>not</u> prepared by EPO or JPO -----	add \$1000/\$500		960/961
2. Search Report was prepared by EPO or JPO -----	add \$860/\$430	+860	970/971

SKIP B, C, D AND E UNLESS country code letters in item 1 are "US", "BR", "BB", "TT", "MX", "IL", "NZ", "IN" or "ZA"

→ <input type="checkbox"/> B. If USPTO did not issue both International Search Report (ISR) and (if box 4(b) above is X'd) the International Examination Report (IPER), -----	add \$1000/\$500	+0	960/961
(X)			
(only) → <input type="checkbox"/> C. If <u>USPTO</u> issued ISR but not IPER (or box 4(a) above is X'd), -----	add \$710/\$355	+0	958/959
(one) →			
(of)			
(these) → <input type="checkbox"/> D. If <u>USPTO</u> issued IPER but IPER Sec. V boxes <u>not all</u> 3 YES, -----	add \$690/\$345	+0	956/957
(4) →			
(boxes)			
→ <input type="checkbox"/> E. If international preliminary examination fee was paid to USPTO and Rules 492(a)(4) and 496(b) <u>satisfied</u> (IPER Sec. V <u>all</u> 3 boxes YES for <u>all</u> claims), -----	add \$100/\$50	+0	962/963

27. **SUBTOTAL =** \$860

28. If Assignment box 19 above is X'd, add Assignment Recording fee of ----\$40 +0 (581)

29. Attached is a check to cover the ----- **TOTAL FEES** \$860

Our Deposit Account No. 03-3975

Our Order No.	009901	0281558
	C#	M#



00909

CHARGE STATEMENT: The Commissioner is hereby authorized to charge any fee specifically authorized hereafter, or any missing or insufficient fee(s) filed, or asserted to be filed, or which should have been filed herewith or concerning any paper filed hereafter, and which may be required under Rules 16-18 and 492 (missing or insufficient fee only) now or hereafter relative to this application and the resulting Official document under Rule 20, or credit any overpayment, to our Account/Order Nos. shown above for which purpose a duplicate copy of this sheet is attached.

This CHARGE STATEMENT **does not authorize** charge of the **issue fee** until/unless an issue fee transmittal form is filed

Pillsbury Winthrop LLP
Intellectual Property Group

By Atty: Donald J. Bjrd

Reg. No. 25323

Sig: [Signature]

Fax: (703) 905-2500
Tel: (703) 905-2018

Atty/Sec: DJB/mhn

NOTE: File in duplicate with 2 postcard receipts (PAT-103) & attachments.

09/889414

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION OF

Inventor(s): RAZA, Ali et al

Filed: Herewith

Title: DRUG COMBINATIONS COMPRISING . . .

July 17, 2001

PRELIMINARY AMENDMENT

Hon. Commissioner of Patents
Washington, D.C. 20231

Sir:

Please amend this application as follows:


IN THE SPECIFICATION:

At the top of the first page, just under the title, insert

☒ --This application is the National Phase of International Application
PCT/GB00/00278 filed February 1, 2000 which designated the U.S.
and that International Application

☒ was ☐ was not published under PCT Article 21(2) in English.--

Respectfully submitted,
PILLSBURY WINTHROP LLP
Intellectual Property Group

By: 
Attorney: Donald J. Bird
Reg. No: 25323
Tel. No.: (202) 905-2018
Fax No.: (202) 905-2500

Atty\Sec. DJB/mhn
1600 Tysons Boulevard

McLean, VA 22102
(703) 905-2000

- 1 -

DRUG COMBINATIONS COMPRISING (E)-7-[4-(4-FLUOROPHENYL)-6-ISOPROPYL-2-[METHYL(METHYLSULFONYL)AMINO]PYRIMIDIN-5-YL](3R,5S)-3,5-DIHYDROXYHEPT-6-ENOIC ACID AND AN INHIBITOR, INDUCER OR SUBSTRATE OF P450 ISOENZYME 3A4

The invention concerns safe non-interacting drug combinations of a 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitor, which is (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof (the Agent) and a drug which is either an inducer, inhibitor or a substrate of cytochrome P450, in particular cytochrome P450 isoenzyme 3A4. Particular combinations are useful in treating hyperlipidaemia in humans who are receiving immunosuppressive chemotherapy. A preferred combination is the Agent and a fibrate drug, the use of such a combination in treating hyperlipidaemia in mammals, and medicaments containing such a combination for use in such treatments.

Hypercholesterolaemia is one of the strongest risk factors for atherosclerosis which is associated with coronary artery disease (including angina pectoris, myocardial infarction and mortality), stroke (including cerebro vascular accident and transient ischaemic attack) and peripheral arterial occlusive disease. Several types of hypercholesterolaemia exist. The magnitude of hypercholesterolaemia may have consequences for the therapy, but in general, any reduction of elevated plasma cholesterol levels is generally accepted to result in an improvement of the risk profile. Dietary improvement and increased exercise are essential first steps and should continue even if drug therapy is instituted, but the therapeutic potential of drug therapy is significantly larger. Several types of drug therapy for hypercholesterolaemia are currently available. Guidelines exist for the treatment of hypercholesterolaemia for example, American Heart Association (AHA) (Anon 1988), Updated Sheffield treatment tables (Heart (1998) 80 Supp.2 S1-S29) and Recommendations of the task force of the European Society of Cardiology Guidelines (Pyorala 1994).

HMG-CoA reductase inhibitors are the most widely used prescription medication for the treatment of hypercholesterolaemia. By inhibiting the rate-controlling step in cholesterol biosynthesis, these agents effectively lower the plasma concentrations of atherogenic particles containing cholesterol such as low-density lipoprotein (LDL-C) and very low-density

FOETTT " 47468860

- 2 -

lipoprotein (VLDL-C). Partial inhibition of hepatic cholesterol synthesis causes up-regulation of hepatic membrane LDL-C receptors which are responsible for the clearance of LDL-C from the circulation. In addition, reduced hepatic synthesis of cholesterol is thought to result in a modest reduction in the secretion of VLDL-C particles by the liver. Clinical trials with

5 certain HMG Co A-reductase inhibitors, such as in the Scandinavian Simvastatin Survival Study, confirm a reduction in cardiovascular morbidity and mortality with such agents, and may even promote regression of atherosclerotic vascular lesions. Various HMG Co A-reductase inhibitors are marketed, and are collectively referred to as 'statins'.

- 10 Despite the impressive benefits of statin therapy, less than optimal therapeutic results may be achieved in some subjects, particularly in the more severe classes of hypercholesterolaemia. This can be due to the occurrence of reversible increases in liver transaminase levels at higher dose levels of statins as well as differences in efficacy between different statins. Clinically important (>3 times upper limit of normal [ULN]) elevations in serum alanine
- 15 aminotransferase [ALT]) have been reported for atorvastatin in 0.8 per cent of patients at low doses of atorvastatin and higher at raised doses (European Summary of Product Characteristics [SmPC] for atorvastatin [Lipitor™]). In all cases the effect is dose-related and reversible. In general it is the incidence of ALT increases which limits dose escalation of statins rather than a limit to further increases in efficacy.

20

- The first generation statins (such as lovastatin, pravastatin and simvastatin - prodrug derivatives of fungal metabolites - and fluvastatin) are categorised in that they achieve only a limited cholesterol lowering affect before the dose administered is limited by elevations in serum ALT. Second generation "superstatins" (such as atorvastatin - synthetic compounds-
- 25 structurally distinct from first generation compounds) inhibitors are categorised in that they lower cholesterol levels to a much higher degree than the earlier first generation of statins before their dose is limited by serum ALT levels. Atorvastatin has been successful over the first generation of statins. Since its launch in the USA atorvastatin has reached sales in 1998, doubling from 1997, of \$2.2 billion, capturing 38% of new prescriptions for cholesterol-
- 30 lowering agents in the US and is now the most widely prescribed hypolipidaemic agent in the US (Warner-Lambert 1998 annual results).

- 3 -

- An additional adverse event, reported for statins in general, is myopathy, defined as symptoms of muscle pain, tenderness and weakness, with creatinine kinase (CK) values $>10 \times$ Upper Limit of Normal (ULN). This adverse event is not considered to be dose related, and in
- 5 addition the adverse events are potentially more serious, and consequently more problematical. In severe cases this can lead to rhabdomyolysis, which is a rare life threatening condition sometimes associated with renal failure. The incidence of raised CK levels ($>10 \times$ ULN - on 2 occasions at least 1 week apart with symptoms = myositis according to FDA) for statins has been reported as 3.1 per cent. (SmPC for atorvastatin). Myopathy and
- 10 rhabdomyolysis have been particularly associated with taking a statin in combination with gemfibrozil, niacin, cyclosporin or erythromycin, (Hunninghake H. Et al. Current Opinion in Lipidolgy (1992), 3, 22-28) which are all substrates for P450 isoenzyme 3A4. The increase in adverse events associated with taking a combination of a statin drug with one of the other drugs mentioned above is probably due to a drug:drug interaction likely related to the
- 15 metabolism of most statins also by the same cytochrome P450 isoenzyme 3A4. Therefore when a drug which is also metabolised by P450 3A4 is administered alongside a statin which also is metabolised by P450 3A4, the side effects discussed above are more likely to occur. Increase in the side effects, such as muscle damage, is thought to be due to elevated statin levels in muscle cells inhibiting farnesylation and geranylgeranylation of muscle proteins .
- 20 Elevated levels of statins may be caused by any drug which affects P450 3A4. Therefore, currently on the labels of all commercially available statins the use of the statin in combination with drugs that are metabolised by P450 3A4 is not recommended and is contraindicated in certain cases.
- 25 Nearly all drugs are metabolised to some degree in the human, generally to a less lipid soluble compound which is more easily excreted by the kidney or in liver bile. The liver is the major site of drug metabolism and many drug metabolising enzymes occur at high concentration in the endoplasmic reticulum (which form microsomes upon homogenisation) of liver parenchymal cells (hepatocytes). Cytochrome P450 represents a major class of drug
- 30 metabolising enzymes and exists as a family of isoenzymes found in hepatic microsomes. Six specific P450 isoenzymes are responsible for the metabolism of most of the commonly used

- 4 -

drugs, namely P450 1A2, 2C9, 2C19, 2D6, 2E1 and 3A4.

A major disadvantage of the currently available "super statin", atorvastatin, is that atorvastatin is metabolised by cytochrome P450 enzymes, in particular 3A4, which may
5 cause drug interactions with other drugs which are inducers, inhibitors or substrates of the same P450 enzyme which metabolises atorvastatin. All of the first generation of statins are metabolised by P450 also. However, the rate of metabolism of pravastatin is sufficiently low that it is considered less susceptible to clinically relevant drug interactions. Therefore despite the lower efficacy of pravastatin, in its currently available doses, at reducing
10 hypercholesterolaemia this is currently the statin of choice in combination with other drugs where the possibility of drug interactions is unacceptably high.

(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]
(3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof (the
15 calcium salt of which is disclosed in Figure 1 below), hereinafter referred to as the Agent, is also a statin and belongs to the class of what is now starting to be called a "superstatin".

The Agent is disclosed in European Patent Application, Publication No. 0521471, and in Bioorganic and Medicinal Chemistry, (1997), 5(2), 437-444 as an inhibitor of HMG-CoA
20 reductase which is a major rate-limiting enzyme in cholesterol biosynthesis. The Agent is described as useful in the treatment of hypercholesterolaemia, hyperlipoproteinaemia and atherosclerosis.

The Agent is not metabolised by cytochrome P450 3A4 and therefore does not possess the
25 same potential for drug interaction shared with the currently available "super statin", i.e. atorvastatin, or any of the other currently available statins.

Therefore we present as a feature of the invention a non-interacting drug combination comprising a HMG CoA reductase inhibitor, which is the Agent, and a drug which is an
30 inhibitor, inducer or substrate of P450 in particular, isoenzyme 3A4.

- 5 -

As a further feature of the invention we present use of a HMG CoA reductase inhibitor, which is the Agent, in the preparation of a medicament for use in combination therapy with a drug which is an inhibitor, inducer or substrate of P450, in particular, isoenzyme 3A4.

- 5 As a further feature of the invention we present use of a drug which is an inhibitor, inducer or substrate of P450, in particular, isoenzyme 3A4 in the preparation of a medicament for use in combination therapy with a HMG CoA reductase inhibitor, which is the Agent.

- As a further feature of the invention we present a pharmaceutical formulation comprising the
10 Agent, a drug which is an inducer, inhibitor or substrate of P450 isoenzyme 3A4 and a pharmaceutically-acceptable diluent, carrier or adjuvant.

- As a further feature of the invention we present a pharmacy pack comprising a first drug which is the Agent and a second drug which is an inducer, inhibitor or substrate of P450
15 isoenzyme 4A4.

- By the term "inducer of P450 " we mean a drug which increases the rate at which a P450 enzyme, in particular isoenzyme 3A4, metabolises a substrate, for example by increasing the activity of the P450 enzyme, decreasing the rate of biological inactivation of the P450 enzyme
20 or by increasing the rate of transcription of the P450 gene.

- By the term "inhibitor of P450" we mean a drug which lowers the rate at which a P450 enzyme, in particular isoenzyme 3A4, metabolises a substrate, for example by lowering the activity of the P450 enzyme or by lowering the rate of transcription of the P450 gene.

25

By the term "substrate of P450" we mean a drug which is metabolised by a P450 enzyme, in particular isoenzyme 3A4.

- By the term "non-interacting drug combination" we mean a drug combination for which there
30 is no adverse affect to the patient by its administration through the mechanism of drug metabolism by cytochrome P450 isoenzyme 3A4. It is recognised that in certain instances a

drug interaction may nevertheless occur between two such drugs when in combination through a completely different mechanism not involving drug metabolism, such as affecting drug absorption.

- 5 Whether a drug is an inhibitor, inducer or substrate of a P450 enzyme can be easily determined by procedures known to the skilled person. Such procedures may involve the exposure of a radiolabelled drug to hepatocytes or hepatocyte microsomes or isolated P450 enzyme and the use of analytic techniques, such as HPLC, in determining metabolite formation. A specific procedure is described herein.

10

By the term "combination" we mean either that the Agent and the drug of the combination are administered together in the same pharmaceutical formulation or that the Agent and the drug are administered separately. When administered separately components of the combination may be administered to the patient simultaneously or sequentially.

15

We have found that the Agent is not metabolised significantly by the major cytochrome P450 isoenzymes 1A2, 2C9, 2C19, 2D6 and 3A4. This is a further feature of the invention.

- Preferred non-interacting combinations of the invention include those in which the Agent is
20 combined with a drug which is also involved in lowering cholesterol and is also an inducer, inhibitor or substrate of P450 3A4. Examples include fibrates, such as bezafibrate, clofibrate, ciprofibrate, fenofibrate and gemfibrozil (preferably fenofibrate), and niacin. Specific embodiments of this preferred feature are described in Section B below.

- 25 Preferred non-interacting combinations of the invention include those in which the Agent is combined with a drug which is involved in treating cardiovascular conditions and which is also an inhibitor, inducer or substrate of P450 3A4. Examples include digitoxin, diltiazem, losartan, nifedipine, quinidine, verapamil and warfarin.

- 30 Preferred non-interacting combinations of the invention include those in which the Agent is combined with cyclosporin and /or tacrolimus (FK506) and therefore has utility in treating

- 7 -

elevated cholesterol levels in patients who are about to, or have recently undergone, a transplantation operation. Specific embodiments of this preferred feature are described below.

Preferred patients in which the combination of the invention is to be administered are those
5 who suffer from myopathy or rhabdomyolysis or who have already been found to suffer from myopathy or rhabdomyolysis when treated with HMG Co A reductase inhibitor which is metabolised by P450 3A4, for example atorvastatin, simvastatin and lovastatin.

Other features of the invention include those described above wherein the Agent is used at
10 doses of 5 to 80mg per day. When a dose range of 5 to 80 mg per day is referred to herein for the Agent other particular dosage ranges, which are further independent aspects of the invention, include (as appropriate) 10 to 80 mg per day, 10 to 60 mg per day, 10 to 40 mg per day, 5 to 40 mg per day, 5 to 20 mg per day, 10 to 20 mg per day, 20 to 60 mg per day, 20 to 40 mg per day and 40 to 60 mg per day. Particular dosages are 5, 10, 20, 40 and 80mg per
15 day. A particularly suitable starting dose of the Agent in the methods referred herein is 5 to 10 mg per day, especially 10 mg per day.

P450 3A4 substrates include; acetaminophen, aldrin, aflentanil, amiodorane, astemizole, benzphetamine, budesonide, carbamazepine, cyclophosphamide, cyclosporin, dapsone,
20 digitoxin, diltiazem, diazepam, erythromycin, etoposide, flutamide, hydroxyarginine, ifosfamide, imipramine, lansoprazole, lidocaine, lovastatin, midazolam, nifedipine, omeprazole, quinidine, rapamycin, retenoic acid, steroids, tacrolimus, teniposide, theophylline, toremifene, triazolam, troleandomycin, verapamil, warfarin, zatosetron and zonisamide.

25

P450 3A4 inhibitors include; clotrimazole, ethinylestradiol, gestodene, itraconazole, ketoconazole, miconazole, diltiazem, naringenin, erythromycin, cyclosporin and triacetyloleandomycin.

- 8 -

P450 3A4 inducers include; carbamazepine, dexamethasone, phenobarbital, phenytoin, rifampin, sulfadimidine, sulfinipyrazone and triacetyloleandomycin.

Examples of other P450 inducers, inhibitors or substrates include those mentioned in

- 5 Drug Metabolism Reviews (1997) Vol 29, Issue 1+2, pages 413-580, Rendic, S. and Di Carlo, F. J. "Human cytochrome P450 enzymes,: A status report summarising their reactions, substrates, inducers and inhibitors".

Dosages of the Agent may be administered according to the cholesterol lowering effect

- 10 desired from a range of 5 to 80 mg per day in any number of unit dosages. Dosages of the drug which is an inducer, inhibitor or substrate of P450 3A4 are those which are advised for each drug, or which are commercially available. Advantageously, due to the lack of interaction at the level of P450 3A4, the skilled person may dose the Agent with a drug which is an inducer, inhibitor or substrate of P450 3A4 with out needing to make any adjustments.

15

The dose ranges and dosages described above are further independent features of the invention.

Preferably the Agent is bis[(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-

- 20 [methyl(methylsulfonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid] calcium salt (illustrated in figure 1).

Experimental

- 25 The experiment below is used to determine the in vitro metabolic fate of [¹⁴C]- labelled Agent in human hepatocytes and, in addition, to determine the specific P450 isozymes involved in [¹⁴C]- labelled Agent metabolism, if any. The latter experiment involves an investigation of the effects of P450 selective chemical inhibitors (see Table 1) on the metabolism of [¹⁴C]- labelled Agent.

30

- 9 -

COMPOUND: [¹⁴C]- labelled Agent.

Chemical name: Bis [(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid] calcium salt

5 **Isomer:** 3R,5S,6E Stereoisomer

Molecular weight: 1001.16 (Ca salt)

Formulation ingredients: The labelled Agent is dissolved in water to produce a solution suitable for addition to the incubates.

10 **TISSUE SOURCE** Human liver, suitable for the preparation of microsomes and hepatocytes, obtained from The International Institute for the Advancement of Medicine (Exton, USA). Human hepatocytes may, in addition, be obtained from Biowhittaker Ltd. or United Kingdom Human Tissue Bank (Leicester, England).

15

EXPERIMENTAL PROCEDURES

(1) METABOLISM OF [¹⁴C]- LABELLED AGENT BY HUMAN HEPATOCYTES

20 [¹⁴C]- labelled Agent (1 μM or higher concentration if required for analytical sensitivity) was incubated with hepatocytes in culture obtained from two human organ donors. Cultures were terminated with ethanol after 0, 6, 24 and 48 hours of incubation and stored at approximately -20°C until analysed. The metabolic competence of the hepatocytes was confirmed at the time of incubation by examining their ability to metabolise [¹⁴C]-ethoxycoumarin (25 μM);

25 aliquots were removed into methanol at the same time points as for the test compound.

Following incubation of [¹⁴C]-ZD4522 with hepatocytes, metabolite profiles were generated by High Performance Liquid Chromatography (HPLC). The ability of hepatocytes to metabolise [¹⁴C]-ethoxycoumarin was confirmed by HPLC.

- 10 -

ASSESSMENT OF DATA

Data generated was assessed with regard to the following:-

- (1) Assess whether human hepatocytes metabolise [^{14}C]- labelled Agent.
- (2) Quantitate the amount of each metabolite formed.

5 (2) ENZYMES INVOLVED IN METABOLISM OF THE AGENT

[^{14}C]- labelled Agent (at an appropriate concentration) was incubated with human hepatic microsomes in the absence and presence of selective P450 inhibitors (see Table 1). Similar incubations of [^{14}C]- labelled Agent with individual heterologously expressed P450 isoenzymes was also performed. Incubations were terminated by the addition of an appropriate organic solvent. Metabolite profiles of the incubates are generated by HPLC.

Table 1 Selective chemical inhibitors of P450 isozymes

P450 isozyme	Selective inhibitor
1A2	Furafylline
2C9	Sulphaphenazole
2C19	Omeprazole
2D6	Quinidine
3A4	Ketoconazole

15 ASSESSMENT OF DATA

Data generated during this study was assessed with regard to the following:-

- (a) The rate and extent of metabolism of [^{14}C]- labelled Agent.
- (b) The ability of the selective P450 inhibitors to decrease the metabolism of [^{14}C]- labelled Agent was compared in order to determine the isozyme(s) involved in the metabolism of [^{14}C]- labelled Agent.

- 11 -

The ability of individual expressed P450 isoforms to metabolise [^{14}C]- labelled Agent was assessed to aid determination of the P450 isozyme(s) involved in the metabolism of [^{14}C]- labelled Agent.

- 5 (c) These *in vitro* data can be used to predict the variability of the pharmacokinetics of the Agent in the population and the likely effects on the pharmacokinetics of the Agent during co-administration with known P450 enzyme inhibitors/inducers.

It was found that the Agent was not significantly metabolised by whole hepatocytes and that this was inhibited by sulphaphenazole and omeprazole.

10

FOR TREATING HYPERLIPIDAEMIA AND ASSOCIATED CONDITIONS IN POST TRANSPLANT PATIENTS RECEIVING IMMUNOSUPPRESSIVE THERAPY.

- 15 Two common drugs used in suppressing the human immune system, cyclosporin and tacrolimus (formerly called FK506), are known to be metabolised by cytochrome P450 3A4.

In particular cyclosporin is also a known inhibitor of P450 3A4 and is therefore likely to reduce the metabolism of any other drug which is metabolised by P450 3A4.

- 20 Therefore where immunosuppressive therapy is prescribed, such as with the drugs cyclosporin and tacrolimus (especially cyclosporin), the attendant physician must be cautious as to any other therapy which may be jointly presented to the patient in combination.

- Immunosuppressive therapy is most commonly used before, during and after human transplant operations. In particular with cardiac transplants the attendant physician may wish to also place the patient on statin drug therapy to reduce future incidents of coronary heart disease, stroke, peripheral arterial occlusive disease or peripheral vascular disease, particularly
- 25 in patients with elevated cholesterol or in normolipidaemic patients with other risk factors associated with heart disease. In particular within this special patient group (human transplant patients), the patients are at high risk of developing accelerated atherosclerosis in the transplant organ in an aggressive fashion and within a short period of time due, in part, to the surgical damage to the blood vessels during transplantation, any previously underlying
- 30 untreated conditions and the immunosuppressive therapy. Hyperlipidaemia is common after transplantation even in patients who did not suffer hyperlipidaemia prior to transplantation, incidence 60-80% of recipients.

- 12 -

It is known that certain immunosuppressive drugs, such as steroids, cyclosporin and tacrolimus, raise cholesterol levels in patients (Wierzbicki AS (1999) IJCP 53 (1) 54-59). In addition cyclosporin and tacrolimus may raise the levels of fibrinogen and lipoprotein (a) in the patient, further accelerating the progression of atherosclerosis in the transplant patient

5 (Hohaye H, Clin.Transplant (1997) 11, 225-230 and Hilbrands LB, J.Am.Soc.Nephrol (1995) 5, 2073-2081). This unusually accelerated atherosclerosis is present in about 20% of heart transplant patients at 1 year and 40-65% at 5 years (Chang G. Et al. American Heart Journal (1998), 136(2), 329-334). The incidence of accelerated atherosclerosis has been reported as causing a 1-18% incidence of CHD at one year and 20-50% incidence of CHD at 3 years in

10 cardiac transplant patients (Erdoes LS, J.Vasc.Surg. (1995) 22, 434-440). Lovastatin, pravastatin and simvastatin have all shown to lower cholesterol levels in heart transplant patients. In a placebo controlled study pravastatin increased survival of transplant patients by 1 year and significantly reduced the incidence of haemodynamic organ rejection. Because of the lower incidence of serious drug interaction with the immunosuppressive therapy pravastatin

15 is currently the statin drug of choice in post transplant treatment regimes. However, as discussed above, pravastatin does not lower lipid/cholesterol levels to such a great extent as, for example, atorvastatin.

We have discovered that the Agent is extremely effective at treating hypercholesterolaemia in

20 patients following transplantation and that the Agent is not metabolised by cytochrome P450 isoenzyme 3A4. Therefore we have found through the use of the Agent in a clinical study that the Agent may be conveniently dosed to patients who are undertaking immunosuppressive therapy without any clinically significant side effects associated with the concomitant dosing of the Agent and the immunosuppressive drug(s) and, in addition, also achieve much higher

25 levels of cholesterol lowering than has previously been achieved, such as by the use of pravastatin.

We present as the first feature of the invention a method of providing safe non-interacting cholesterol lowering therapy to a human patient undertaking immunosuppressive

30 chemotherapy which method comprises administering to the patient the Agent.

- 13 -

Particular patients undertaking immunosuppressive chemotherapy who may benefit from the method of the invention are those who:

- 1) suffer primary (type IIa) hypercholesterolaemia ($\text{LDL-L} \geq 135$ and $\text{TG} < 200$);
- 2) suffer combined (type IIb) hypercholesterolaemia ($\text{LDL-C} \geq 135$ and $\text{TG} \geq 200$);
- 3) patients with established CHD or other atherosclerotic disease, such as PVD, stroke or peripheral arterial occlusive disease;
- 4) patients who are at high risk of developing CHD or other atherosclerotic disease, such as described above, because of a combination of risk factors. The term "high risk" is defined in the "Recommendations of Second Joint Task Force of European and other Societies on Coronary Prevention", (Wood, D. et. al. European Heart Journal, Atherosclerosis and Journal of Hypertension 1998) as absolute CHD risk of $\geq 20\%$ over 10 years or will exceed 20% if projected to age 60 years. Whether a patient is at high risk or not may be determined by the charts which accompany the above recommendations and which charts are incorporated herein by reference. For example a male patient in his 40s who smokes and has a systolic blood pressure of 180 mm Hg or higher and a total plasma cholesterol concentration of 7 mmol/L or higher will be classified as high risk. Similarly other guidelines for reducing risk factors may be applied such as those described in;

a) JAMA, June 16, 1993-Vol 629, No.23, Pages 3015-3023 - "Summary of the NCEP Adult Treatment Panel II Report" - specifically Figure 1. Page 3018-3019 which is incorporated herein by reference.

b) Post Graduate Medical Journal 1993; 69(811): 359-369 - "Management of hyperlipidaemia: guidelines of the British Hyperlipidaemic Association"- specifically Table V and Table VI which are incorporated herein by reference.

- 14 -

c) Heart 1998; 80 Supplement 2:S1-S29 - "Joint British recommendations on prevention of coronary heart disease in clinical practice" - specifically Figure 1 on pages S4-S5, which is incorporated herein by reference.

d) The Lancet 1995; December 2, Vol.346, 1467-1471 - "Sheffield risk and treatment table for cholesterol lowering for primary prevention of coronary heart disease" - specifically the Table appearing at page 1468, which is incorporated herein by reference.

5) patients who suffer type I or II diabetes;

6) patients who are about to or have already undertaken a heart transplant;

The statin therapy may be administered so as to achieve in the patient undertaking immunosuppressive chemotherapy.

1) A reduction in the internal thickness of coronary artery atheroma of $\geq 30\%$ as measured by IVUS.

2) A reduction of LDL-C of at least 30, 40, 50%.

3) A maintenance or increase of HDL-C of at least 5, 10, 15%.

4) A change in any of the above values better than pravastatin at a similar dose and over the same period.

As a further feature of the invention, and due also to the fact that the Agent is not metabolised to any significant extent by P450 isoenzymes, it is possible to administer, more safely than before, to a patient receiving immunosuppressive therapy a fibrate and the Agent. As discussed earlier the administration of a fibrate and a statin has previously been associated with a higher incidence of rhabdomyolysis and myopathy. In addition fibrate drugs do interact with cyclosporin due to both being metabolised by the same P450 isoenzyme. Therefore, the use of

a statin and a fibrate drug in combination with immunosuppressive therapy was previously contraindicated due to the likelihood of possible serious interactions (Hunninghake 1992, Wanner C. Kidney Int. (1995) 52(suppl.), S60-S62; and Katznelson S. Contributions Nephrol. (1997) 120, 97-104). However, if possible, it would be advantageous to also administer a

- 5 fibrate alongside a statin since fibrates are known to lower different lipoproteins than statins and therefore their combined pharmacology would be complementary in reducing even further the likelihood of CHD and other diseases mentioned above associated with the formation of atherosclerosis. Therefore the possibility of combining the Agent, which is not metabolised by P450 3A4, with a fibrate and an immunosuppressive therapy offers the additional possibility of
- 10 lowering cholesterol to a greater extent in such patients than previously achieved and more safely than could previously be achieved by the administration of a statin, a fibrate and an immunosuppressive drug.

Fibrate drugs are thought to act through peroxisomal proliferating activator receptor- α

- 15 (PPAR- α) and affect gene activation at a number of genes involved in atheroma. Patients on fibrate drugs show improved LDL subfraction distribution (reduced VLDL and raised HDL), reduced LDL and reduced triglyceride levels, and possible advantages through improving insulin sensitivity. Examples of fibrate drugs include, bezafibrate, ciprofibrate, fenofibrate and gemfibrozol.

20

By use of the term "safe non-interacting statin therapy" we mean that the Agent is not metabolised by P450 3A4 and therefore does not affect the metabolism of the immunosuppressive therapy or *vice versa*.

- 25 Diseases and conditions in which immunosuppressive therapy may be prescribed include, in addition to organ transplantation mentioned above, autoimmune diseases, including rheumatic disorders, such as, rheumatoid arthritis, osteoarthritis, lupus erthematosus; and other autoimmune disorders such as idiopathic thrombocytopenic purpura, autoimmune haemolytic anaemia and acute glomerulonephritis.

30

- 16 -

The agent may be administered at the same time as the immunosuppressive chemotherapy, or if not at the same time within a short time period of administration of the immunosuppressive therapy, such as in the same day, within 6, 3, 2 or 1 hour.

- 5 The Agent may be administered according to the cholesterol lowering effect desired from a range of 5-80 mg per day in any number of unit dosages, preferable once a day dosing. Ideal doses are 10, 20 and 40 mg once per day. Preferred doses are 20 and 40mg once per day.

- Particular immunosuppressive drugs which may be combined with the Agents are those which are metabolised by liver enzymes, such as by P450 3A4, and therefore are not likely to have a drug interaction with the Agent. Examples include those described above, cyclosporin and tacrolimus, as well as corticosteroids, which are also metabolised in the liver. Examples of corticosteroids include prednisone (especially used for organ transplantation). Preferably at least one of the immunosuppressive agents, if more than one agent is used, is either cyclosporin or tacrolimus, preferably cyclosporin.
- 10
15

Example

- The following non-limiting example is of a clinical trial to demonstrate the performance of this aspect of the invention.
- 20

PROTOCOL

Title:

- A Double-blind, Parallel Group Study to Assess the Change in Coronary Artery Atheroma Burden Post Cardiac Transplantation as Measured via IVUS after 12 Months Dosing with the Agent versus Pravastatin
- 25

Objectives:

- The primary objective of the study is to measure change in maximal mean intimal thickness of the anterior descending coronary artery as assessed by intravascular ultrasonography (IVUS) (read centrally) after 12 months of treatment with the Agent or pravastatin. A change from
- 30

- 17 -

baseline of $\geq 30\%$ in intimal thickness is considered clinically significant.

The secondary objectives of the study are to measure the effects on coronary artery atheroma burden and to compare effects of the Agent with the following assessments:

- evidence of organ rejection as assessed by adverse event reports.
- measurement of LDL-C, HDL-C, apoB, apoA-I, Lp (a) concentrations, *ex vivo* platelet aggregation, fibrinogen, PAI-I, and the concentrations of circulating markers of vascular inflammation.
- comparison of lipid values after 52 weeks of treatment.
- measurement of inflammatory markers after 52 weeks of treatment (HLA antigen VCAM/ICAM expression as assessed by biopsy).
- to determine the drug's safety and tolerability.

Type and number of subjects: Approximately 40 men and women (aged 18 years and older) post cardiac transplant with hypercholesterolemia and triglycerides < 400 mg/dl at the time of randomisation.

Trial treatment: Once daily doses of the Agent (10 mg) or pravastatin (10 mg) for two weeks, then titration of dose to 20 mg of the Agent or pravastatin 20 mg. After 4 weeks the dose should be titrated up to 40 mg of the Agent or 40 mg pravastatin. Patients who have had their dose titrated up to 40 mg may have their dose titrated down to 20 mg, at the discretion of the investigator.

Duration of treatment: Eligible subjects randomised to 1 of 2 treatment groups, the Agent or pravastatin, for 52 weeks.

- 18 -

Primary measure:

Mean change from baseline in maximal mean intimal thickness, as assessed by IVUS (read centrally).

Secondary measures:

Percent change from baseline in LDL-C at 6 and 12 months.

5

Percent change from baseline in total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), LDL-C/HDL-C, TC/HDL-C, non-HDL-C/HDL-C, and triglycerides (TG).

10

Percent change from baseline in ApoB, ApoB/ApoA-1, ApoA-1, Lp (a), and particle subfractions at 6 and 12 months.

Percentage of subjects on each of the possible titrated doses at 12 months.

Endocardial rejection will be considered an adverse event.

15

Percent change from baseline in inflammatory markers (HLA antigen level and ICAM/VCAM expression).

Safety evaluation as determined by adverse events, physical examination, and laboratory data.

TRIAL DESIGN

- 20 This is a multicenter, randomized, double-blind, parallel-group clinical trial. Within 1 to 4 weeks post surgery, subjects are randomized to receive either the Agent or pravastatin for 52 weeks. Subjects start treatment at a dose of 10 mg of either the Agent or pravastatin at Visit 2 and the dose is titrated to 20 mg at Visit 3 during the forced titration period. At Visit 4 and subsequent visits, the investigator has the option to increase each drug up to 40 mg during
- 25 the optional titration period. Patients who have had their dose titrated up to 40 mg may have their dose titrated back down to 20 mg at the investigator's discretion.

TRIAL DESIGN

	Pre-transplant	Forced Titration		Optional Titration							
Visit	1	2	3	4	5	6	7	8	9	10	11
Week (W)/		W0	W2	W4	M2	M3	M4	M5	M6	M9	M12
Month (M)											
Agent (mg)		10	20	≥ 20*							
PRAVASTATIN (mg)		10	20	≥ 20*							
Randomisation**											

* Subjects who are tolerating 20 mg of the Agent or Pravastatin at Visit 4 may have their dose titrated up to 40 mg, at the discretion of the investigator.

** Subjects should be randomized within 4 weeks of cardiac transplantation and must not have received any other lipid lowering therapy post-surgery.

Inclusion criteria

- 5 (1) have undergone cardiac transplantation up to four weeks prior to randomization
- (2) fasting TG concentrations of <4.52 mmol/L (400 mg/dl)

Exclusion criteria

- 10 Any of the following is regarded as a criterion for exclusion from the trial:
 - (1) Use of other cholesterol lowering drugs or lipid lowering dietary supplements or food additives post-transplantation prior to entering the study
 - (2) history of serious or hypersensitivity reactions to other HMG-CoA reductase inhibitors
- 15 (3) pregnant women, women who are breast feeding, and women of child bearing potential who are not using chemical or mechanical contraception or have positive serum pregnancy test (a serum β -Human chorionic gonadotropin [β -HCG] analysis)

- 20 -

- (4) Subjects with a history of diabetic ketoacidosis within the past 5 years are excluded.
- (5) uncontrolled hypothyroidism defined as a thyroid stimulating hormone (TSH) >1.5 times the ULN at Visit 2 or subjects whose thyroid replacement therapy was initiated within the last three months
- 5 (6) use of concomitant medications as detailed below - except immune suppressants and diazepam
- (7) current alcohol and/or drug abuse
- (8) active liver disease or hepatic dysfunction as defined by elevations of ≥ 1.5 times the ULN at Visit 2 in any of the following liver function tests: ALT, AST, or bilirubin
- 10 (9) serum CK > 3 times ULN at Visit 2
- (10) serum creatinine > 220 $\mu\text{mol/L}$ (2.5 mg/dl)
- (11) subjects with cancer or with a history of cancer who, in the opinion of the investigator, have more than a minimal chance of recurrence
- (12) participation in another investigational drug trial less than 4 weeks before
15 randomization into the trial
- (13) subjects randomized to double-blind treatment who subsequently withdrew cannot re-enter this trial
- (14) serious or unstable medical or psychological conditions that, in the opinion of the investigator, would compromise the subject's safety or successful participation in the
20 trial
- (15) subjects taking cyclic hormone replacement therapy (HRT), cyclic oral contraceptive therapy (OCT), a depot progesterone injection, or subjects whose non-cyclic HRT or OCT was initiated within the last 3 months

DISALLOWED MEDICATIONS

CLASS OF DRUG	GENERIC NAME
Antibiotics/ antifungals	Erythromycin Base Erythromycin Ethyl Succinate, Acetyl Sulfisoxazole Rifampicin Fluconazole Ketaconazole Itraconazole
Anti-epileptics/ antidepressants	Phenytoin Phenobarbitol Fluoxetine Carbamazepine
Acne treatment	Isotretinoin
Antiulcer drugs	Cimetidine Cisapride

T.O.T.T. 446860

CLASS OF DRUG	GENERIC NAME
Systemic Steroids	Triamcinolone Acetonide
	Triamcinolone Diacetate
	Betamethasone
	Sodium Phosphate
	Betamethasone Acetate
	Hydrocortisone
	Hydrocortisone Acetate
	Hydrocortisone Sodium Phosphate
	Hydrocortisone Sodium Succinate
	Cortisone Acetate
	Dexamethasone
	Dexamethasone Acetate
	Dexamethasone Sodium
	Prednisone
	Methylprednisolone
	Methylprednisolone Acetate
	Methylprednisolone Sodium
	Succinate
	Prednisolone Tebutate
	Prednisolone Sodium Phosphate
	Methyltestosterone
	Fluoxymesterone
Antihistamine	Astemizole
	Terfenadine

CLASS OF DRUG	GENERIC NAME
Lipid Regulation	Niacin/Nicotinic Acid Probucol Psyllium Preparations Clofibrate Cholestyramine Colestipol Hydrochloride Gemfibrozil Atorvastatin Lovastatin Pravastatin (except study medication) Simvastatin Fluvastatin Cerevastatin Fish oils (any dose) lipid lowering dietary supplements lipid lowering food additives
Hormone Therapy	Estrogen and progesterone combinations which are bi or tri phasic.

Friedewald Equation

The LDL-C level is calculated from the Friedewald equation as follows:

5 For SI units (mmol/l)

$$\text{LDL-C} = \text{Total cholesterol} - [\text{HDL-C} + \text{Triglycerides}/2.2]$$

For non-SI units (mg/dl):

$$\text{LDL-C} = \text{Total cholesterol} - [\text{HDL-C} + \text{triglycerides}/5]$$

Summary of NCEP Goals for Lipid Management^a

NCEP Risk Category	Target LDL-C (NCEP)
No CHD/PVD and 1 or no risk factors	< 160 mg/dL
No CHD/PVD and 2 or more risk factors	< 130 mg/dL
Clinically evident CHD/PVD	≤ 100 mg/dL

- 5 ^a Second Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Bethesda (MD): National Institutes of Health, National Heart and Lung Institute 1993 Sep Report No.: 93-3095.
NCEP National Cholesterol Education Program.

10

FOR TREATING HYPERLIPIDAEMIA, AND ASSOCIATED CONDITIONS, USING A COMBINATION OF THE AGENT AND A FIBRATE DRUG OR NIACIN

- 15 Myopathy and rhabdomyolysis have been associated with taking a statin in combination with gemfibrozil, niacin, cyclosporin or erythromycin, (HMG CoA reductase inhibitors, Hunninghake, Current Opinion in Lipidology (1992) 3, 22-28) which are all substrates for P450 3A4. Additionally, adverse events associated with taking a fibrate drug have also been reported to increase with concomitant statin therapy, such as a myositis-like syndrome, which occasionally occurs in patients receiving gemfibrozil, increases to 5% of patients when
20 a statin is also administered.

- Combination of a statin with a fibrate drug is contraindicated on the labels, both in the USA and Europe, of all commercially available statins. Despite the possibility of the occurrence of serious drug interactions doctors do prescribe combination therapy of a statin and a fibrate
25 drug to patients with more severe levels of hypercholesterolaemia, such as in patients with

- 25 -

5 familial combined hyperlipidaemia, where the risk of a serious drug interaction is outweighed by the benefits of the combination therapy. It is recommended that where combination therapy of a fibrate drug and a statin is prescribed that patients should have their CK value determined on a regular basis, typically every 6-weeks, until a stable pattern is established. Therapy is stopped if muscle symptoms occur in association with elevated CK activity. However, as quoted from the US label of Lipitor™ "there is no assurance that such monitoring [of CK levels] will prevent the occurrence of severe myopathy".

10 We have discovered that the Agent is extremely effective at treating mixed hyperlipidaemia and hypertriglyceridaemia in patients when combined with a fibrate drug and that the Agent is not metabolised by cytochrome P450 isoenzyme 3A4. Therefore we have found through the use of the Agent in a clinical study that the Agent may be conveniently dosed to patients who are also taking a fibrate drug without any clinically significant side effects associated with the concomitant dosing of the Agent and the fibrate drug. In addition much higher levels of lipid
15 lowering than has previously been achieved can be achieved by the use of the Agent and a fibrate drug. The combination is of most use in mixed hyperlipidemia where the LDL and VLDL and TGs are all elevated.

20 We present as the first feature of the invention a method of providing safe non-interacting lipid lowering combination therapy to a mammal, including a human patient, preferably a patient suffering mixed hyperlipidaemia and hypertriglyceridaemia, which method comprises administering to the patient the Agent and a fibrate drug or niacin.

25 By the term "combination" as used herein we mean either (1) that the Agent and the fibrate drug of the combination are administered together in the same pharmaceutical formulation or (2) that the Agent and the drug are administered separately. When administered separately components of the combination may be administered to the patient simultaneously or sequentially.

30 By the term "fibrate drug" we mean the class of drugs which are based around the structure/activity of fibric acid and such drugs include the following commercially available

- 26 -

versions; bezafibrate, clofibrate, ciprofibrate, fenofibrate and gemfibrizol, preferably fenofibrate.

Preferred patients in which the combination of the invention is to be administered are those
5 who have already been found to suffer from myopathy or rhabdomyolysis when treated with a statin and/or with a fibrate drug which is metabolised by P450 3A4.

Particular patients who may benefit from the method of the invention are those who:

- 10 1) suffer combined (type IIb) hypercholesterolaemia (typically LDL-C \geq 135 mg/dL and TG \geq 200 mg/dL);
- 2) suffer familial (type IV and V) hypercholesterolaemia;
- 3) patients suffering secondary hypercholesterolaemia from such conditions as:
 - 15 a) diabetes (type I or II),
 - b) nephrotic syndrome,
 - c) uremia,
 - d) hyperthyroidism, and
 - e) obstructive liver disease.
- 4) patients with established CHD or other atherosclerotic disease, such a PVD, stroke or
20 peripheral arterial occlusive disease;
- 5) patients who are at high risk of developing CHD or other atherosclerotic disease,⁵ such as described above, because of a combination of risk factors. The term "high risk" is defined in the "Recommendations of Second Joint Task Force of European and other Societies on Coronary Prevention", (Wood, D. et. al. European Heart Journal, Atherosclerosis and
25 Journal of Hypertension 1998) as absolute CHD risk of \geq 20% over 10 years or will exceed 20% if projected to age 60 years. Whether a patient is at high risk or not may be determined by the charts which accompany the above recommendations and which charts are incorporated herein by reference. For example a male patient in his 40s who smokes and has a systolic
30 blood pressure of 180 mm Hg or higher and a total plasma cholesterol concentration of 7 mmol/L or higher will be classified as high risk. Similarly other guidelines for reducing risk factors may be applied such as those described in;

- 27 -

a) JAMA, June 16, 1993-Vol 629, No.23, Pages 3015-3023 - "Summary of the NCEP Adult Treatment Panel II Report" - specifically Figure 1. Page 3018-3019, which is incorporated herein by reference.

b) Post Graduate Medical Journal 1993; 69(811): 359-369 - "Management of hyperlipidaemia: guidelines of the British Hyperlipidaemic Association"- specifically Table V and Table VI, which are incorporated herein by reference.

c) Heart 1998; 80 Supplement 2:S1-S29 - "Joint British recommendations on prevention of coronary heart disease in clinical practice" - specifically Figure 1 on pages S4-S5, which is incorporated herein by reference.

d) The Lancet 1995; December 2, Vol.346, 1467-1471 - "Sheffield risk and treatment table for cholesterol lowering for primary prevention of coronary heart disease" - specifically the Table appearing at page 1468, which is incorporated herein by reference.

15 The statin therapy may be administered so as to achieve in the patient receiving a fibrate drug or niacin:

- 1) a reduction of LDL-C of at least 30, 40, 50, 60, 70 or 80%.
- 2) a maintenance or increase of HDL-C of at least 5, 10, 15%.
- 3) a reduction in triglycerides of at least 10, 20, 30 or 40%.

The combination of the fibrate, or niacin, and the Agent may be applied as separate dosage forms, which may be taken simultaneously or sequentially, or in a combined dosage form.

The combination of the fibrate and the Agent will also have an additive or synergistic effect on the reduction in LDL-C, maintenance or increase of HDL-C or reduction in triglyceride in the patients blood.

In addition the combination of niacin and the Agent may be applied as separate dosage forms, which may be taken simultaneously or sequentially, or in a combined dosage form. The

combination of the fibrate and the Agent will also have an additive or synergistic effect on

- 28 -

the reduction in LDL-C, maintenance or increase of HDL-C or reduction in triglyceride in the patients blood.

Doses of the Agent which are administered are at the discretion of the attendant physician

- 5 generally taking into account the severity of the disease, the age, weight and sex of the patient. However typical doses will be from 5 to 80 mg per day orally, preferably as a once a day oral tablet form.

- 10 Doses of the fibrate drug or niacin which are administered in the combination of the invention also are at the discretion of the attendant physician taking into account all of the above factors plus in particular which drug is used.

For clofibrate (such as Atromid-S®) 20-30 mg/kg body weight daily in 2 or 3 divided oral doses after meals is typical.

15

For bezofibrate (such as Bezalip®) 400 mg once a day orally, after food at night or in the morning, is typical.

- 20 For fenofibrate (such as Lipantil®) 200 mg once a day, or 62 mg three times a day, with food is typical.

For gemfibrozil (such as Lopid®) 600 mg twice a day orally is typical.

For ciprofibrate (such as Modalim®) 100 mg once a day orally is typical.

25

For niacin (NIASPAN®, an extended release niacin formulation, and preferred feature) 500mg once to four times daily, preferably twice or four times daily.

A preferred fibrate drug is fenofibrate.

30

Preferably the AGENT is administered to a patient receiving niacin at 10mg or 40mg daily doses.

The particular aspect of this invention is illustrated by the following non-limiting examples:

5 Clinical Trial

A Randomised, Non-controlled, Single-centre, Open-label, 3-way Crossover Trial to Assess the Effect of Co-administration of the Agent and Fenofibrate on the Pharmacokinetics of Each Compound in Healthy Male Volunteers

Objectives: The primary objective of this trial is to assess the effect of co-administration of the Agent and fenofibrate on the pharmacokinetics of both the Agent and fenofibrate

The safety of all volunteers will be ensured by clinical monitoring

Type and number of volunteers: 14 healthy male volunteers

Trial design: The trial will be a randomised, non-controlled, 3-way crossover study carried out at a single centre

Trial treatment: This trial will consist of three 7-day treatment periods (Periods A, B, and C). Volunteers will receive, in random order, a 10 mg capsule of the Agent once daily for 7 days, a 67 mg fenofibrate capsule 3 times daily for 7 days and the combination for 7 days.

There will be a minimum of a 3-week washout between each trial period.

Duration of treatment: The study will consist of 3 periods of 7-day dosing (a total of 21 dosing days) with a 3-week washout between dosing in Periods A, B and C.

Primary endpoints: The primary endpoints are:

- AUC(0-24) and C_{max} of the Agent in the presence and absence of fenofibrate
- AUC(0-8) and C_{max} of fenofibrate in the presence and absence of the Agent

- 30 -

**Secondary
endpoints:**

the secondary endpoints are:

- t_{\max} , $t_{1/2}$, C_{\min} for the Agent in the presence and absence of fenofibrate
- t_{\max} , $t_{1/2}$, C_{\min} for fenofibrate in the presence and absence of the Agent
- safety assessments: symptoms, blood pressure and pulse rate, ECG, clinical chemistry, haematology and urinalysis

T.C.T. 446880

TRIAL PLAN

Summary of procedures - overall plan for Trial Periods A, B and C

Trial Days	Medical	Doses of the Agent / fenofibrate or combination	P & BP	12 lead ECG	Safety Blood & Urine	Kinetics of the Agent	Kinetics Fenofibrate
Pre-trial	+		+	+	+ ^a		
-1					+ ^b		
1		+				+ ^c	+ ^c
2		+			+ ^b	+ ^d	+ ^c
3		+				+ ^d	+ ^c
4		+			+ ^b		
5		+					
6		+			+ ^b	+ ^d	+ ^c
7		+	+	+		+ ^d	+ ^c
8					+ ^b	+ ^d	
9						+ ^d	
10					+ ^b	+ ^d	
Post-trial	+		+	+	+ ^a	+ ^d	

^aFull clinical chemistry, haematology and urine labstix.

^bClinical chemistry only: urea, creatinine, total protein, albumin, uric acid, total bilirubin (and unconjugated and conjugated bilirubin if total bilirubin raised), alkaline phosphatase, alanine aminotransferase (ALT), aspartate aminotransferase (AST), gamma glutamyltransferase, creatine kinase (CK), sodium, potassium, calcium, cholesterol and triglycerides.

^cPre-dose all trial periods.

^dOnly trial periods when volunteers receive the Agent

10 Only trial periods when volunteers receive fenofibrate

P = pulse, BP = blood pressure

- 32 -

TRIAL PLAN II**Trial Day 7 in Periods A, B and C**

Time	P & BP (L)	12 lead ECG	Safety blood & urine ^c	Kinetics of the Agent ^b	Kinetics fenofibrate ^c	Meals & Fluids
Pre-dose	+	+		+	+	B
Dose (0 h)						D
0.5 h				+	+	
1 h				+	+	
2 h				+	+	D
3 h	+	+		+	+	
4 h				+	+	M, F
5 h	+	+		+	+	
6 h				+	+	
8 h				+	+	S
10 h				+		M
12 h	+	+		+		F
14 h						S
18 h				+		W
24 h	+	+	+ ^a	+		
30 h				+		
48 h				+		
54 h				+		
72 h			+ ^a	+		

^aclinical chemistry only: urea, creatinine, total protein, albumin, uric acid, total bilirubin (and unconjugated and conjugated bilirubin if total bilirubin raised), alkaline phosphatase, ALT, AST, gamma glutamyltransferase, CK, sodium, potassium, calcium, cholesterol and triglycerides.

^bOnly trial periods when volunteers receive the Agent

^cOnly trial periods when volunteers receive fenofibrate

10 L = lying; P = pulse; BP = blood pressure; D = drink; S = snack; M = meal; F = free access to permitted fluid and food; W = free access to water only

1 OBJECTIVES

Primary objective

The primary objective of this trial is to assess the effect of co-administration of the Agent and fenofibrate on the pharmacokinetics of both the Agent and fenofibrate.

5 Secondary objective

There is no secondary objective for this trial.

The safety of all volunteers will be ensured by clinical monitoring.

Design

10 The trial will be a randomised, non-controlled, open-label, 3-way crossover study carried out at a single centre.

Volunteers will receive 3 treatment regimens in random order:

- 10 mg of the Agent once daily for 7 days
- fenofibrate (Lipantil™) 67 mg x 3 daily for 7 days
- the Agent (10 mg once daily) and fenofibrate (Lipantil™, 67 mg x 3 daily) given in
15 combination for 7 days

There will be a minimum of 3 weeks (21 days) washout between each treatment period.

Inclusion criteria

For inclusion in the trial, volunteers must meet all of the following criteria:

- male, aged between 18 and 65 years inclusive
- 20 • normal clinical examination, including medical history, resting electrocardiogram (ECG) and 24-hour continuous ambulatory ECG (if not performed in the past 12 months)
- negative screens for serum hepatitis B surface antigen and hepatitis C antibody and a normal screen for ferritin within the previous 12 months
- 25 • weight not differing by more than 20% from the desirable weight (Metropolitan Height and Weight Tables)

Exclusion criteria

Volunteers must be excluded from the trial if any of the following criteria are met:

- use of any medication or therapy, including drugs of abuse
- receipt of another new chemical entity in the 4 months before dosing in this trial (a
5 new chemical entity is defined as a compound which has not been submitted for
marketing authorisation)
- participation in another trial within 3 months before the start of the present trial, apart
from non-invasive methodology trials in which no drugs were given
- any acute illness within 2 weeks before the start of the trial
- 10 • any clinically significant abnormalities in clinical chemistry, haematology or
urinalysis results. In addition the following clinical chemistry parameters should be
no greater than the upper limit of normal: total bilirubin, ALT, AST and CK
- risk (in the investigator's opinion) of transmitting, through blood or other body
15 fluids, the agents responsible for acquired immune deficiency syndrome (AIDS),
hepatitis B or hepatitis C
- definite or suspected personal history or family history of adverse drug reactions, or
hypersensitivity to drugs with a similar chemical structure to the Agent or related
statins, or fenofibrate and related fibrate drugs
- history or presence of gastrointestinal, hepatic, biliary or renal disease or other
20 condition known to interfere with absorption, distribution, metabolism or excretion
of drugs
- history of Gilbert's syndrome
- if participation in the trial would result in the volunteer donating more than 1350 ml
of blood in the 12 months before the end of the trial
- 25 • excessive intake of alcohol, defined as a maximum weekly intake of greater than
28 units (1 unit equals half a pint of beer or a measure of spirits)

- 35 -

- treatment in the previous 3 months with any drug known to have a well-defined potential for hepatotoxicity (eg, halothane)
- clinical judgement by the investigator or the volunteer's general practitioner that the volunteer should not participate in the trial

5 Volunteer restrictions

Volunteers will be required to:

- abstain from taking any medication (including over-the-counter remedies) from 96 hours before Trial Day 1 to 72 hours after receiving the last dose of the Agent or morning dose of fenofibrate in each trial period unless the investigator has given prior consent
- fast from midnight on the night before each trial day and eat a light breakfast on arrival on Trial Day 1 to 7 in each trial period
- refrain from driving, cycling, using machinery (drills, sanders, sharp instruments etc.) for 24 hours after receiving first dose on Trial Day 7 in each period
- remain for 24 hours after receiving first dose on Trial Day 7 in each trial period
- abstain from smoking, consuming grapefruit, grapefruit juice, liquorice or caffeine-containing drinks or foods (eg, coffee, tea, cocoa, chocolate and cola) from midnight before Trial Day 1 until 72 hours after receiving the last dose of the Agent or morning dose of fenofibrate in each trial period
- abstain from drinking alcohol from 96 hours before Trial Day 1 until 72 hours after receiving the last dose of the Agent or morning dose fenofibrate in each trial period
- refrain from physical exercise from 96 hours before Trial Day 1 until 72 hours after receiving the last of the Agent or morning dose of fenofibrate in each trial period
- refrain from potentially hazardous work or activities, from receiving the first dose of the Agent or fenofibrate until the post-trial medical

- 36 -

- abstain from donating blood during the trial and for 3 months following their last dose of trial treatment

Formulation, presentation and storage

Dosage and administration

- 5 Capsules of the Agent or fenofibrate will be taken orally with 200 ml of purified water with the volunteer sitting in an upright position.

On Trial Days 1 to 7 of each treatment period, volunteers will receive one of the following treatments:

- 1 x 10 mg capsule of the Agent to be taken between 08:30 and 09:30 hours
- 10 • 3 x 67 mg fenofibrate capsules
 - the 1st capsule to be taken between 08:30 and 09:30 hours
 - the 2nd capsule to be taken between 16:30 and 17:30 hours with food
 - the 3rd capsule to be taken between 22:30 and 23:30 hours with food-
- 1 x 10 mg capsule of the Agent and 3 x 67 mg fenofibrate capsules:
 - 15 – 1 capsule of the Agent and the 1st fenofibrate capsule to be taken simultaneously between 08:30 and 09:30 hours
 - the 2nd fenofibrate capsule to be taken between 16:30 and 17:30 hours with food
 - the 3rd fenofibrate capsule to be taken between 22:30 and 23:30 hours with food
- 20

On Trial Days 1 to 6 of each trial period, volunteers will visit the unit daily and will be allowed to leave the unit immediately after administration of doses of the Agent, fenofibrate or the the Agent / fenofibrate combination, except on Trial Day 7 when volunteers will remain resident for 24 hours.

25

In trial periods when the volunteers are randomised to fenofibrate, they will take the further 2 doses of fenofibrate at home. The volunteers will be provided with 1 pot of fenofibrate to be

- 37 -

taken as outlined above. Volunteers will be issued with a pre-set timer to ensure that the dose is taken at the correct time, and a diary card to document the dose was taken.

When the the Agent and fenofibrate are given to the volunteers, the tear-off labels will be attached to the appropriate case report form (CRF). The investigator must ensure that each

5 volunteer receives the correct treatment.

Clinical and laboratory assessments

Primary endpoints

The following parameters will be measured as primary endpoints:

- AUC(0-24) and C_{max} of the Agent in the presence and absence of fenofibrate
- 10 • AUC(0-8) and C_{max} of fenofibrate in the presence and absence of the Agent

Secondary endpoints

The following parameters will be measured as secondary endpoints:

- t_{max} , $t_{1/2}$, C_{min} for the Agent in the presence and absence of fenofibrate
- t_{max} , $t_{1/2}$ and C_{min} for fenofibrate in the presence and absence of the Agent
- 15 safety assessments: symptoms, blood pressure and pulse rate, ECG, clinical chemistry, haematology and urinalysis.

Pharmaceutical compositions

- 20 The following Example illustrates, but is not intended to limit, pharmaceutical dosage forms which are suitable for use in the invention as defined herein:

Capsule	mg
The Agent	5.0
25 Lactose	42.5
Cornstarch	20.0
Microcrystalline cellulose	32.0
Pregelatinised starch	3.3

- 38 -

Hydrotalcite 1.1
magnesium stearate 1.1


Capsules containing 1, 2.5 or 10mg of the Agent may be obtained similarly using more or less
5 lactose as appropriate, to maintain a total fill weight of 105mg.

ABBREVIATIONS AND CONVENTIONS USED

Abbreviation	Term
ALT	alanine aminotransferase
ALP	alkaline phosphatase
apo B	apolipoprotein B 100
AST	aspartate aminotransferase
AUC	area under the concentration curve from zero to infinity
AUC(0-t)	area under the curve of plasma concentration against time from zero to time of last quantifiable concentration
CABG	Coronary artery bypass graft
C _{max}	maximum concentration
CK	creatinine kinase
CVA	cerebrovascular accident
ECG	electrocardiogram
EAS	European Atherosclerosis Society
EDTA	ethylenediamine-tetraacetic acid
XGT	Gemma glutaryl transferase
HMG-CoA	3-hydroxy-3-methylglutaryl coenzyme A

Abbreviation	Term
HDL	high-density lipoprotein
HPLC	high-performance liquid chromatography
HRT	hormone replacement therapy
IU	International Units
IVUS	Intravascular ultrasenography
LDL	low density lipoprotein
LDL-C	low density lipoprotein cholesterol
MVA	mevalonic acid
NC	not calculable
NCEP	national cholesterol eduction program
NDSR	national data system for research
THC	tetrahydrocannabinol
TG	triglyceride
$t_{1/2}$	terminal elimination half-life
t_{max}	time of maximum concentration
TC	total cholesterol
TG	triglycerides
TIA	transient ischemic attack
TSH	thyroid stimulating hormone
ULN	upper limit of normal
VLDL	very low-density lipoprotein

- 40 -

Figure 1

Formula I

Claims

1. A non-interacting drug combination comprising a HMG-CoA reductase inhibitor, which is (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof and
5 a drug which is an inhibitor, inducer or substrate of P450 isoenzyme 3A4.
2. A non-interacting drug combination, as claimed in claim 1, wherein the second drug is an inhibitor or inducer of P450 isoenzyme 3A4.
10
3. A non-interacting drug combination, as claimed in either claim 1 or claim 2, wherein each drug is administered together or each drug is administered sequentially.
4. A non-interacting drug combination, as claimed in any claim from 1 to 3, wherein the
15 second drug is used to lower cholesterol and is an inducer, inhibitor or substrate of P450 isoenzyme 3A4.
5. A non-interacting drug combination, as claimed in claim 4, wherein the second drug is selected from bezafibrate, clofibrate, fenofibrate, gemfibrozol and niacin.
20
6. A non-interacting drug combination, as claimed in claim 5, wherein the second drug is fenofibrate.
7. A non-interacting drug combination, as claimed in any claim from 1 to 3, wherein the
25 second drug is used in treating cardiovascular conditions and is also an inhibitor, inducer or substrate of P450 isoenzyme 3A4.
8. A non-interacting drug combination, as claimed in claim 7, wherein the second drug is selected from digitoxin, diltiazem, losartan, nifedipine, quinidine, verapamil and warfarin.

9. A non-interacting drug combination, as claimed in any claim from 1 to 3, wherein the second drug is used in immunosuppression therapy and is an inducer, inhibitor or substrate of P450 isoenzyme 3A4.
- 5 10. A non-interacting drug combination, as claimed in claim 9, wherein the second drug is selected from cyclosporin, tacrolimus and a corticosteroid.
11. A non-interacting drug combination, as claimed in any claim from 1 to 10, wherein (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]
- 10 (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof is dosed at 5, 10, 20, 40 or 80mg once per day.
12. A pharmaceutical formulation comprising (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a
- 15 pharmaceutically acceptable salt thereof, a drug which is an inducer, inhibitor or substrate of P450 isoenzyme 3A4 and a pharmaceutically-acceptable diluent, carrier or adjuvant.
13. A pharmaceutical formulation, as claimed in claim 12, wherein the second drug is a substrate of P450 isoenzyme 3A4 and is selected from acetaminophen, aldrin, alfentanil,
- 20 amiodorane, astemizole, benzphetamine, budenoside, carbamazepine, cyclophosphamide, cyclosporin, dapsone, digitoxin, diltiazem, diazepam, erythromycin, etoposide, flutamide, hydroxyarginine, ifosfamide, imipramine, lansoprazole, lidocaine, lovastatin, losartan, lovastatin, midazolam, nifedipine, omeprazole, quinidine, rapamycin, retenoic acid, steroids, tacrolimus, teniposide, theophylline, toremifene, triazolam, troleandomycin, verapamil,
- 25 warfarin, zatosetron and zonisamide.
14. A pharmaceutical formulation, as claimed in claim 12, wherein the second drug is an inhibitor of P450 isoenzyme 3A4 and is selected from clotrimazole, ethinylestradiol, gestodene, itraconazole, ketoconazole, miconazole, diltiazem, naringenin, erythromycin,
- 30 cyclosporin and triacetyloleandomycin.

15. A pharmaceutical formulation, as claimed in claim 12, wherein the second drug is an inducer of P450 isoenzyme 3A4 is selected carbamazepine, dexamethasone, phenobarbital, phenytoin, rifampin, sulfadimidine, sulfinipyrazone and triacetyloleandomycin.

5

16. A pharmacy pack comprising a first drug which is (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof and a second drug which is an inducer, inhibitor or substrate of P450 isoenzyme 4A4.

10

17. A pharmacy pack, as claimed in claim 16, wherein the second drug is used to lower cholesterol and is an inducer, inhibitor or substrate of P450 isoenzyme 3A4.

18. A pharmacy pack, as claimed in claim 17, wherein the second drug is selected from bezafibrate, clofibrate, fenofibrate, gemfibrozol and niacin.

19. A pharmacy pack, as claimed in claim 16, wherein the second drug is used in treating cardiovascular conditions and is also an inhibitor, inducer or substrate of P450 isoenzyme 3A4.

20

20. A pharmacy pack, as claimed in claim 19, wherein the second drug is selected from, digitoxin, diltiazem, losartan, nifedipine, quinidine, verapamil and warfarin.

21. A pharmacy pack, as claimed in claim 16, wherein the second drug is a substrate of P450 isoenzyme 3A4 and is selected from acetaminophen, aldrin, aflentanil, amiodorane, astemizole, benzphetamine, budesonide, carbamazepine, cyclophosphamide, cyclosporin, dapson, digitoxin, diltiazem, diazepam, erythromycin, etoposide, flutamide, hydroxyarginine, ifosphamide, imipramine, lansoprazole, lidocaine, lovastatin, losartan, lovastatin, midazolam, nifedipine, omeprazole, quinidine, rapamycin, retenoic acid, steroids, tacrolimus, teniposide, theophylline, toremifene, triazolam, troleandomycin, verapamil, warfarin, zatosetron and zonisamide.

30

09889414-111304
PCT/GB00/00278

22. A pharmacy pack, as claimed in claim 16, wherein the second drug is an inhibitor of P450 isoenzyme 3A4 and is selected from clotrimazole, ethinylestradiol, gestodene, itraconazole, ketoconazole, miconazole, diltiazem, naringenin, erythromycin, cyclosporin and
5 triacetyloleandomycin.

23. A pharmacy pack, as claimed in claim 16, wherein the second drug is an inhibitor of P450 isoenzyme 3A4 and is selected from carbamazepine, dexamethasone, phenobarbital, phenytoin, rifampin, sulfadimidine, sulfinpyrazone and triacetyloleandomycin.

10

24. Use of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof in the preparation of a medicament for use in combination therapy with a second drug which is an inducer, inhibitor or substrate of P450
15 isoenzyme 3A4.

25. Use of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof in the preparation of a medicament for use in
20 cholesterol lowering therapy in combination therapy with a second drug which is an inducer, inhibitor or substrate of P450 isoenzyme 3A4.

26. Use, as claimed in claim 25, wherein the second drug is selected from bezafibrate, clofibrate, fenofibrate, gemfibrozil and niacin.

25

27. Use of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof in the preparation of a medicament for use in the treatment of cardiovascular condition in combination with a second which is an inducer,
30 inhibitor or substrate of P450 isoenzyme 3A4.

- 45 -

28. Use, as claimed in claim 27, wherein the second drug is selected from digitoxin, diltiazam, losartan, nifedipine, quinidine, verapamil and warfarin.

29. Use of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-

5 [methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof in the preparation of a medicament for use in cholesterol lowering therapy in a patient receiving immunosuppressive therapy.

30. Use, as claimed in claim 29, wherein the immunosuppressive therapy comprises the

10 administration of a drug selected from cyclosporin, tacrolimus and a corticosteroid.

31. Use of (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-

[methyl(methylsulfonyl)amino]pyrimidin-5-yl] (3R,5S)-3,5-dihydroxyhept-6-enoic acid or a pharmaceutically acceptable salt thereof in the preparation of a medicament for use in

15 cholesterol lowering therapy in combination with a second drug which is selected from bezafibrate, clofibrate, fenofibrate, gemfibrozil and niacin in a patient receiving immunosuppressive therapy.

32 Use as claimed in claim 31 wherein the immunosuppressive therapy comprises the

20 administration of a drug selected from cyclosporin, tacrolimus and a corticosteroid.

FOR UTILITY/DESIGN
CIP/PTCT NATIONAL/PLANT
ORIGINAL/SUBSTITUTE/SUPPLEMENTAL
DECLARATIONS

RULE 63 (37 C.F.R. 1.63)
DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

PW
FORM

Z70475/UST

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name, and I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the **INVENTION ENTITLED: DRUG COMBINATIONS COMPRISING (E)-7-[4-(4-FLUOROPHENYL)-6-ISOPROPYL-2-METHYL (METHYLSULPHONYL)AMINO]PYRIMIDIN-5-YL(3R,5S)-3,5-DIHYDROXYHEPT-6-ENOIC ACID AND AN INHIBITOR, INDUCER OR SUBSTRATE OF P450 ISOENZYME 3A4**

the specification of which (CHECK applicable BOX(ES))
X ☐ A ☐ is attached hereto.
BOX(ES) ☒ B ☐ was filed on _____ as U S Application No. _____

☒ C ☐ was filed as PCT International Application No. PCT/GB00/00278 on 01 February 2000 _____ and (if applicable to U.S. or PCT application) was amended on _____. I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. 1.56. Except as noted below, I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate or 365(a) of any PCT International Application which designated at least one other country than the United States, listed below and have also identified below any foreign application for patent or inventor's certificate, or PCT International Application, filed by me or my assignee disclosing the subject matter claimed in this application and having a filing date (1) before that of the application on which priority is claimed, or (2) if no priority claimed, before the filing date of this application.

PRIOR FOREIGN APPLICATION(S) Number	Country	Date first laid- open or Published	Date Patented or Granted	Priority NOT Claimed
9902593.4	United Kingdom	06.02.1999		
9921063.5	United Kingdom	08.09.1999		
9921064.3	United Kingdom	08.09.1999		

Except as noted below, I hereby claim domestic priority benefit under 35 U.S.C. 119(e) or 120 and/or 365(c) of the indicated United States applications listed below and PCT international applications listed above or below and, if this is a continuation-in-part (CIP) application, insofar as the subject matter disclosed and claimed in this application is in addition to that disclosed in such prior applications, I acknowledge the duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. 1.56 which became available between the filing date of each such prior application and the national or PCT international filing date of this application:

PRIOR U.S. PROVISIONAL, NON PROVISIONAL AND/OR PCT APPLICATION(S) Application No. (series code/serial no.)	Date/MONTH/Year Filed	Status Pending, abandoned, patented	Priority NOT Claimed
---------------------------------------------------------------------------------------------------------------	-----------------------	----------------------------------------	----------------------

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

And I hereby appoint Pillsbury Winthrop LLP, Intellectual Property Group, telephone number (202)861-3000 (to whom all communications are to be directed), and persons of that firm who are associated with USPTO Customer No 909 (see below label) individually and collectively my attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith and with the resulting patent, and I hereby authorize them to delete from that Customer No names of persons no longer with their firm, to add new persons of their firm to that Customer No., and to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct the above Firm and/or an attorney of that Firm in writing to the contrary.

00909

(1) INVENTOR'S SIGNATURE

Name	JOHN	First	S	Middle Initial	PEARS	Family Name
Residence	Macclesfield	City	Cheshire	State/Foreign Country	United Kingdom	Country of Citizenship
Mailing Address	Alderley Park, Macclesfield, Cheshire SK10 4TG, United Kingdom					
(include Zip Code)						

(2) INVENTOR'S SIGNATURE

Name	ALI	First	RAZA	Middle Initial	Family Name	
Residence	Wilmington	City	Delaware	State/Foreign Country	USA	Country of Citizenship
Mailing Address	1800 Concord Pike, PO Box 15437, Wilmington, Delaware 19850-5437, USA					
(include Zip Code)						

☒ OR ADDITIONAL INVENTORS see attached page.

☐ See additional foreign priorities on attached page (incorporated herein by reference).

Atty. Dkt. No. P (M#)

DECLARATION AND POWER OF ATTORNEY

(continued)
Additional Inventors

Page 2
Z70475/UST

(3) INVENTOR'S SIGNATURE

Date: 4 September 2001

Name	HOWARD	First	G	Middle Initial	HUTCHINSON	Family Name
Residence	Wilmington	City	Delaware	State/Foreign Country	USA	Country of Citizenship
Mailing Address	1800 Concord Pike, PO Box 15437, Wilmington, Delaware 19850-5437, USA					
(include Zip Code)						

(4) INVENTOR'S SIGNATURE

Date: 4 September 2001

Name	DENNIS	First		Middle Initial	SCHNECK	Family Name
Residence	Wilmington	City	Delaware	State/Foreign Country	USA	Country of Citizenship
Mailing Address	1800 Concord Pike, PO Box 15437, Wilmington, Delaware 19850-5437, USA					
(include Zip Code)						

(5) INVENTOR'S SIGNATURE

Date: 18th September 2001

Name	TAKAHIKO	First		Middle Initial	BABA	Family Name
Residence	Toyonaka-shi	City	Osaka	State/Foreign Country	Japan	Country of Citizenship
Mailing Address	Shionogi & Co., 1-1, Futabacho 3-chome, Toyonaka-shi, Osaka 561-0825, Japan					
(include Zip Code)						

(6) INVENTOR'S SIGNATURE

Date: 18th September 2001

Name	AKIRA	First		Middle Initial	TOUCHI	Family Name
Residence	Toyonaka-shi	City	Osaka	State/Foreign Country	Japan	Country of Citizenship
Mailing Address	Shionogi & Co., 1-1, Futabacho 3-chome, Toyonaka-shi, Osaka 561-0825, Japan					
(include Zip Code)						

(7) INVENTOR'S SIGNATURE

Date: 18th Sep. 2001

Name	YOSHITAKA	First		Middle Initial	YAMAGUCHI	Family Name
Residence	Toyonaka-shi	City	Osaka	State/Foreign Country	Japan	Country of Citizenship
Mailing Address	Shionogi & Co., 1-1, Futabacho 3-chome, Toyonaka-shi, Osaka 561-0825, Japan					
(include Zip Code)						

(8) INVENTOR'S SIGNATURE

Date:

Name		First		Middle Initial		Family Name
Residence		City		State/Foreign Country		Country of Citizenship
Mailing Address						
(include Zip Code)						

(9) INVENTOR'S SIGNATURE

Date:

Name		First		Middle Initial		Family Name
Residence		City		State/Foreign Country		Country of Citizenship
Mailing Address						
(include Zip Code)						